



# Supporting Information

Pascal et al. 10.1073/pnas.1108073108



**Fig. S1.** (A) Equilibrium molecular dynamics (MD) structure of the 10 nanotubes considered in this study. Each structure is the last snapshot of a 5-ns constant temperature and constant pressure MD run with a finite nanotube. Water molecules outside the nanotube are not shown for clarity. (B) Schematic of the starting MD system configuration for infinite nanotube simulations. The equilibrium structures in A are inserted into a box of 14,000 preequilibrated simple point charge-extended waters (shown in gray dots). All nanotubes are 10-nm long.



**Fig. S2.** Number of enclosed water molecules in finite open-ended carbon nanotube (CNT); (A) 0.8-nm (6,6) CNT; (B) 2.2-nm (16,16) CNT over 5 ns of constant temperature and constant pressure (NPT) dynamics. Convergence is seen after 1 ns of dynamics. The snapshot with number of enclosed waters closest to the average during the last 4 ns (Table S1) was used to form an infinite nanotube and solvated with simple point charge-extended waters (10 Å buffer in the x and y dimension) for long-term NPT dynamics.






Fig. S5. Comparison of the (A) free energy  $\Delta A^0$ , (B) enthalpy  $\Delta U^0$ , and (C) entropy  $\Delta S^0$  of water confined in 10 carbon nanotubes (CNTs) with the simple point charge-extended (SPC-E) water model (this study, red squares) and with the unoptimized parameters of the AMBER96 force field for carbon and the TIP3P water model (oxygen-carbon parameters of  $\epsilon_{C-O} = 0.392$  kJ/mol and  $\sigma_{C-O} = 3.275$  Å, blue filled circles) (1). The good agreement indicates that the results presented in this study are fairly insensitive to the choice of interaction potentials. As a figure of merit, we calculate the free energy of water inside the 0.8-nm (6,6) CNT to be  $-3.98$  kJ/mol using the two-phase thermodynamics method, which is in good agreement with the value of  $-3.34$  kJ/mol in ref. 1 from potential of mean force calculations.

1 Hummer G, Rasaiah JC, Noworyta JP (2001) Water conduction through the hydrophobic channel of a carbon nanotube. *Nature* 414:188–190.

**Table S1. Description and results of finite molecular dynamics (MD) simulations**

CNT	Initial unit cell dimensions, Å		No. water molecules in unit cell	No. enclosed waters	
	x/y	z		avg	±
6,6	28.14	127.17	9,393	36	2
7,7	29.50	127.17	12,949	102	2
8,8	30.85	127.17	14,618	172	3
9,9	32.20	127.17	16,253	219	4
10,10	33.56	127.17	17,923	294	6
11,11	34.92	127.17	19,624	369	5
12,12	36.28	127.17	21,174	465	5
14,14	39.00	127.17	24,499	673	6
16,16	41.70	127.17	27,978	913	7
20,20	47.13	127.17	34,969	1,536	7

Results are obtained from the last 4 ns of a 5-ns constant temperature and constant pressure run. The number of enclosed waters was obtained by counting all water molecules within a cylindrical region defined by the carbon nanotube (CNT) end caps and the CNT radius.

**Table S2. Density of water molecules inside carbon nanotubes (CNTs) during the last 4 ns of 5-ns constant temperature and constant pressure dynamics**

CNT	CNT diameter, nm		Total density, g/mL		Effective density, g/mL	
	avg	±	avg	±	avg	±
6,6	0.8256	0.0008	0.202	0.013	0.496	0.048
7,7	0.9612	0.0006	0.425	0.010	0.887	0.039
8,8	1.0954	0.0006	0.552	0.004	1.034	0.027
9,9	1.2300	0.0006	0.562	0.006	0.963	0.014
10,10	1.3664	0.0008	0.605	0.012	0.983	0.018
11,11	1.5020	0.0008	0.627	0.008	0.972	0.020
12,12	1.6368	0.0366	0.667	0.007	0.990	0.011
14,14	1.9094	0.0008	0.708	0.006	0.989	0.012
16,16	2.1802	0.0010	0.738	0.006	0.983	0.011
20,20	2.7248	0.0010	0.795	0.004	0.994	0.007

The total density assumes an internal volume defined by the entire CNT diameter. The effective density has an effective radius which accounts for the carbon van der Waals radii. Convergence to the bulk density begins to be seen approximately after the (10,10) CNT.